

The Joint Position Statement on Adult and Adolescent Sexual Assault Patients in the Emergency Care Setting was posted previously on the ENA and IAFN Websites. It is being published in the *Journal of Emergency Nurses* (ENA) and the *Journal of Forensic Nursing* (JFN) in order to reach the widest possible audience.

Adult and Adolescent Sexual Assault Patients in the Emergency Care Setting

Emergency Nurses Association | International Association of Forensic Nurses

Description

Patients seeking care in the emergency setting after sexual assault are an at-risk, vulnerable population. It is imperative that they have access to and receive patient-centered and trauma-informed care that addresses their medicolegal and psychosocial needs (Plichta, Clements, & Houseman, 2007). Patient-centered care is respectful of and responsive to individual patient preferences, needs and values, ensuring that patient choices guide all clinical decisions (Committee on Quality of Health Care in America, 2001). Trauma-informed care involves understanding the connection between presenting symptoms and behaviors, on the one hand, and the individual's past trauma history, on the other (Hodas, 2006).

Comprehensive care for these patients requires extensive planning, education and expertise on the part of emergency care providers and their collaborative partners (Sampsel, Szobota, Joyce, Graham, & Pickett, 2009). The emergency nurse is a key member of this team and instrumental in coordinating access to appropriate healthcare providers, collaborating with community-based victim advocates, social workers, and law enforcement personnel (Cybulska, 2013; Linden, 2011). Many nurses have education and certification in the expanded practice role of sexual assault nurse examiner (SANE) in order to provide care in an empowering setting, different from traditional medical care that often left patients feeling re-victimized (Cybulska, 2013; Linden, 2011). The SANE completes a time sensitive examination that includes: assessing, treating and documenting injuries, identifying risks and providing preventative treatment for negative health outcomes associated with sexual assault including exposure to infection,

unintended pregnancy, and long term psychological and physical sequelae, collecting evidence, maintaining the chain of custody, safety planning, and providing support with appropriate community referrals (Plichta et al., 2007; Sampsel et al., 2009; Cybulska, 2013). Immediate medical and psychological care directly impacts the patient's well-being and contributes to the beginning stages of the healing process (Plichta et al., 2007; Cybulska, 2013; Linden, 2011). Successful physical and emotional outcomes, as well as potential prosecution of sexual offenders, require emergency care settings to be prepared to provide competent care and referrals for the sexual assault patient (Plichta et al., 2007).

ENA/IAFN Position

It is the position of the Emergency Nurses Association and the International Association of Forensic Nurses that:

1. Sexual assault patients are provided a safe and private environment upon arrival in an emergency care setting, with access to a community-based advocate at any time during their stay.
2. Emergency nurses use a trauma-informed approach throughout the sexual assault patient's complex plan of care.
3. Sexual assault patients receive consistent, objective, immediate medical care, as well as options for the collection of evidence by emergency nurses and physicians knowledgeable of jurisdictional guidelines and protocols for evidence collection.
4. Whenever possible, forensic nurses with specific training as sexual assault nurse examiners are consulted or assigned to care for this patient population.
5. Sexual assault patients receive medically appropriate sexually transmitted disease prophylaxis and emergency contraception without barriers, according to recommended Centers for Disease Control and American

Copyright © 2017 Emergency Nurses Association. "Adult and Adolescent Sexual Assault Patients in the Emergency Care Setting" is jointly published by the Emergency Nurses Association, the *Journal of Emergency Nurses* published by Elsevier Inc. and the International Association of Forensic Nurses, the *Journal of Forensic Nursing* published by Wolters Kluwer. All rights reserved.

Congress of Obstetricians and Gynecologists guidelines, respectively.

6. Emergency nurses receive continuing education on medical and forensic sexual assault evaluation and maintain access to current legislative guidelines and protocols for proper examination and reporting options.
7. Emergency nurses collaborate with multiple agencies to develop an individualized, multidisciplinary approach to treatment, evaluation and continuity of care to minimize the patient's short and long term physical and psychological trauma.
8. Emergency nurses participate in community education and research to identify and implement best practice standards of care for the sexual assault patient.
9. Healthcare facilities recognize that they have an obligation to provide appropriate medical forensic intervention when a sexual assault patient presents for care, whether or not the facility has a SANE program.
10. Healthcare facilities are cognizant of jurisdictional laws regarding all aspects of the sexual assault medical/forensic exam.
11. Healthcare facilities support SANE practitioners by developing, sustaining or maintaining their own programs, or establishing relationships with other facilities.
12. Lead trauma centers develop and maintain SANE services to address the health needs of this patient population.

Background

Sexual assault patients regularly present to the emergency care setting for treatment following their traumatic event (Plichta et al., 2007). In the past, they have often been treated by emergency department personnel who lacked training in medical forensic evidence collection, and those with training often did not perform exams frequently enough to maintain proficiency and competency (Littel, 2001). The result was poor documentation and improper evidence collection. Historically, emergency physicians have found it difficult to dedicate the amount of time required to provide the necessary level of care for this population while still managing the other patients in the emergency department (Plichta et al., 2007). The understanding that these patients require complex clinical management in a setting that often has significant time constraints, led to the use of specially trained examiners to provide care for sexual assault patients (Sampsel et al., 2009; Linden, 2011; Houmes, Fagan, & Quintana, 2003). One study found proper documentation in 100% of SANE documented evidence kits compared to 79% in non-SANE kits. It also found that proper evidence collection specimens were reported in 96% of SANE collected kits compared to 86% when collected by physicians (Houmes et al., 2003). Case law has repeatedly found

SANE-collected kits to be among the strongest courtroom evidence in supporting victim testimony, and SANE nurses themselves have been noted to be very credible witnesses (Plichta et al., 2007; Linden, 2011; Houmes et al., 2003). The role of specially trained sexual assault nurse examiners has been supported by the American College of Emergency Physicians (ACEP) and they are the FBI's preferred examiners for victims and suspected perpetrators of criminal sexual acts (Linden, 2011; Houmes et al., 2003).

Guidelines for the treatment of patients after a sexual assault have been issued by the Department of Justice, ACEP, the American Congress of Obstetricians and Gynecologists and the World Health Organization (Linden, 2011; Houmes et al., 2003). The Centers for Disease Control and Prevention continually update the recommendations for pharmacological treatment after exposure to potential sexually transmitted diseases including HIV. Across these organizations it is recommended that patients presenting to an emergency care setting be assessed for acute traumatic physical injuries and offered forensic evidence collection according to jurisdictional protocols (Linden, 2011). To reduce further re-victimization by providing prompt care and overall better services, patients should be referred to clinicians with education and experience in systematically managing this population (Houmes et al., 2003; Campbell, 2006). Using specialized examiners has alleviated previous issues of emergency department increased wait times, poor clinical outcomes and patient dissatisfaction (Plichta et al., 2007; Girardin, 2005). The new standard of care includes deliberate and timely crisis intervention, proper medical care and evidence collection, and complete coordination and follow up with members of the community sexual assault response team (Plichta et al., 2007; Girardin, 2005). Additionally, the presence of a community-based advocate during the examination as a source of support for the patient improves both medical service and legal system responses for victims (Campbell, 2006).

Resources

American Congress of Obstetrics and Gynecologists: Emergency Contraception: Resource Overview. Retrieved from <http://www.acog.org/Womens-Health/Emergency-Contraception>

Centers for Disease Control and Prevention: 2015 Sexually Transmitted Diseases Treatment Guidelines: <http://www.cdc.gov/std/tg2015/>

Sexual Assault Forensic Examiner Technical Assistance: <http://www.safeta.org/>

Forensic Nursing Education Guidelines: <http://www.forensicnurses.org/?page=EducationGuidelines>

Position Statement: Collaboration With Victim Advocates: http://www.forensicnurses.org/resource/resmgr/Position_Papers/IAFN_Position_Statement-Advo.pdf

Position Statement: The Use of Emergency Contraception
Post Sexual Assault Statement: http://www.forensicnurses.org/resource/resmgr/Position_Papers/IAFN_Position_Statement-Emer.pdf

Position Statement: DNA Evidence Collection from the Oral Cavity: http://www.forensicnurses.org/resource/resmgr/Position_Papers/DNA_Evidence_Collection_From.pdf

Management of the Patient with the Complaint of Sexual Assault: <http://www.acep.org/content.aspx?id=29562>

A National Protocol for Sexual Assault Medical Forensic Examinations Adults/Adolescents: <https://www.ncjrs.gov/pdffiles1/ovw/241903.pdf>

Guidelines for medico-legal care for victims of sexual violence: <http://apps.who.int/iris/bitstream/10665/42788/1/924154628X.pdf>

Committee on Health Care for Underserved Women Opinion on Sexual Assault: <http://www.acog.org/Resources-And-Publications/Committee-Opinions/Committee-on-Health-Care-for-Underserved-Women/Sexual-Assault>

References

- Campbell, R. (2006). Rape survivors' experiences with the legal and medical systems: do rape victim advocates make a difference? *Violence Against Women*, 12(1), 30–45.
- Committee on Quality of Health Care in America. (2001). *Crossing the quality chasm: A new health system for the 21st century*. Washington, D.C.: National Academy Press.
- Cybulska, B. (2013). Immediate medical care after sexual assault. *Best Practice & Research Clinical Obstetrics and Gynecology*, 27(1), 141–149.
- Girardin, B. (2005). The sexual assault nurse examiner: a win-win solution. *Topics in Emergency Medicine*, 27(2), 124–131.
- Hodas, G. (2006). Responding to childhood trauma: the promise and practice of trauma informed care. *National Association of State Mental Health Program Directors (NASMHPD)*.
- Houmes, B. V., Fagan, M. M., & Quintana, N. M. (2003). Establishing a sexual assault nurse examiner (SANE) program in the emergency department. *The Journal of Emergency Medicine*, 25(1), 111–121.
- Linden, J. A. (2011). Clinical practice. Care of the adult patient after sexual assault. *The New England Journal of Medicine*, 365(9), 834–841.
- Littel, K. (2001). Sexual assault nurse examiner programs: Improving the community response to sexual assault victims. *Office for Victims of Crime Bulletin*, 4, 1–19.
- Plichta, S. B., Clements, P. T., & Houseman, C. (2007). Why SANEs matter: models of care for sexual violence victims in the emergency department. *Journal of Forensic Nursing*, 3(1), 15–23.
- Sampsel, K., Szobota, L., Joyce, D., Graham, K., & Pickett, W. (2009). The impact of a sexual assault/domestic violence program on ED care. *Journal of Emergency Nursing*, 35(4), 282–289.

Authors

Authored by
Katie Bush, MA, RN, CEN, SANE-A

Reviewed by

ENA 2016 Position Statement Committee Members

E. Marie Wilson, MPA, RN, Chairperson

Melanie Crowley, MSN, RN, CEN

Kathy Dolan, MSHA, RN, CEN, CPHRM

Ellen Encapera, RN, CEN

Elizabeth Stone, MSN, RN, CPEN

Justin Winger, PhD, MA, BA, RN

IAFN Chief Executive Officer, Interim

Jennifer Pierce-Weeks, RN, SANE-A, SANE-P

IAFN 2016 Board of Directors Liaison

Jana French, MS, BSN, RN, SANE-A,

Board of Directors

ENA 2016 Board of Directors Liaison

Sally Snow, BSN, RN, CPEN, FAEN

ENA 2016 Staff Liaison

Dale Wallerich, MBA, BSN, RN

Developed: 2016.

Approved by the ENA Board of Directors:

July 2016.

Approved by the IAFN Board of Directors:

August 2016

©Emergency Nurses Association, 2016.

©International Association of Forensic Nurses, 2016

This position statement, including the information and recommendations set forth herein, reflects ENA and IAFN's current position with respect to the subject matter discussed herein based on current knowledge at the time of publication. This position statement is only current as of its publication date, and is subject to change without notice as new information and advances emerge. The positions, information and recommendations discussed herein are not codified into law or regulations. In addition, variations in practice, which take into account the needs of the individual patient and the resources and limitations unique to the institution, may warrant approaches, treatments and/or procedures that differ from the recommendations outlined in this position statement. Therefore, this position statement should not be construed as dictating an exclusive course of management, treatment or care, nor does adherence to this position statement guarantee a particular outcome. ENA and IAFN's position statements are never intended to replace a practitioner's best nursing judgment based on the clinical circumstances of a particular patient or patient population. Position statements are published by ENA and IAFN for educational and informational purposes only, and ENA and IAFN do not "approve" or "endorse" any specific sources of information referenced herein. ENA and IAFN assume no liability for any injury and/or damage to persons or property arising out of or related to the use of or reliance on any position statement.